

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.

1. REPORT DATE (DD-MM-YYYY) 15-02-2012			2. REPORT TYPE Conference Proceedings		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE A Multi-Scale Model of the Turkish Straits System					5a. CONTRACT NUMBER	
					5b. GRANT NUMBER	
					5c. PROGRAM ELEMENT NUMBER 0602435N	
6. AUTHOR(S) Cheryl Ann Blain, Mustafa Kemal Cambazoglu, Ewa Jarosz					5d. PROJECT NUMBER	
					5e. TASK NUMBER	
					5f. WORK UNIT NUMBER 73-6263-09-5	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004					8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7320-09-0094	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research One Liberty Center 875 North Randolph Street, Suite 1425 Arlington, VA 22203-1995					10. SPONSOR/MONITOR'S ACRONYM(S) ONR	
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution is unlimited.						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT A multi-scale coastal ocean model representing the entire Turkish Straits System ( Aegean-Marmara-Black Seas and connecting straits) is one-way coupled to a 1 km basin-scale ocean model of the same region. The modeled three-dimensional circulation and density structure of the TSS is examined through comparison to observations of currents and density taken during the TSS08 sea trial.						
15. SUBJECT TERMS coastal models, circulation, bosporus, dardanelles, Marmara Sea						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT		18. NUMBER OF PAGES	
a. REPORT	b. ABSTRACT	c. THIS PAGE	UU		1	
Unclassified	Unclassified	Unclassified				
19a. NAME OF RESPONSIBLE PERSON Cheryl Ann Blain					19b. TELEPHONE NUMBER (Include area code) 228-688-5450	

20120217340

# A MULTI-SCALE MODEL OF THE TURKISH STRAITS SYSTEM

Cheryl Ann Blain <sup>1\*</sup>, Mustafa Kemal Cambazoglu <sup>2</sup> and Ewa Jarosz <sup>1</sup>

<sup>1</sup> Oceanography Division, Naval Research Laboratory, Stennis Space Center, MS, USA - cheryl.ann.blain@nrlssc.navy.mil

<sup>2</sup> Dept. Marine Science, University of Southern Mississippi, Stennis Space Center, MS, USA

## Abstract

A multi-scale coastal ocean model representing the entire Turkish Straits System (Aegean-Marmara-Black Seas and connecting straits) is one-way coupled to a 1 km basin-scale ocean model of the same region. The modeled three-dimensional circulation and density structure of the TSS is examined through comparison to observations of currents and density taken during the TSS08 sea trial.

**Keywords:** Coastal Models, Circulation, Bosphorus, Dardanelles, Marmara Sea

Two narrow, shallow straits, i.e. the Dardanelles and the Bosphorus, form a physical connection between the Marmara Sea and its adjacent water bodies, the Aegean Sea to the southwest and the Black Sea to the northeast. This collection of seas and straits is known as the Turkish Strait System (TSS). Saline, dense water from the Aegean flows in a deep, lower layer through the Marmara Sea to the Black Sea while fresher, lighter Black Sea water flows in a surface layer to the Aegean Sea. Though the TSS dynamics are the result of interconnections between the interconnected straits and ocean basins, earlier modeling efforts (e.g., Staneva et al., 2001; Oguz, 2005; Kourafalou and Tsirlas, 2007; Kanarska and Maderich, 2008) have focused on dynamical studies of individual straits or seas. Often the geometric complexity, broad range of spatial scales present, and computational requirements to represent such disparity have prevented study of the TSS as a whole.

For this study, we utilize state-of-the-art modeling practices to capture the range of spatial scales, geometric complexity and interconnected dynamics of the TSS (Figure 1). A model based on unstructured grids has the resolution, using a minimum element edge length of 20 m, necessary to model flow in the narrow straits whose minimum width is approximately 600 m. The Advanced Circulation Model (ADCIRC), solves the three-dimensional flow and transport equations using a finite element discretization with a terrain-following, generalized, stretched coordinate system applied in the vertical (Luettich and Westerink, 2004; Dresback and Kolar, 2009). Flexibility of the finite element mesh not only captures the fine scales within the straits but is also able to represent mesoscale variability in the Marmara Sea while coupling to a basin scale model in the Aegean and Black Seas. Basin-wide dynamics are captured by the HYbrid Coordinate Ocean Model, HYCOM, which applies the finite difference method over a structured grid to solve the primitive mass and momentum balance equations (Bleck, 2002). HYCOM's hybrid vertical coordinate allows the use of three vertical coordinate types (depth, terrain-following and isopycnal) which better represents thermohaline dynamics in waters of rapidly varying bathymetric change. Within the HYCOM Aegean-Marmara-Black Sea model (HYCOM-AMB), both straits are represented as idealized channels since the current resolution (~1.3 km) is not sufficient to resolve the geometry of the straits.



Fig. 1. ADCIRC finite element model domain for the Turkish Strait System including a zoom of the mesh refinement in the Dardanelles Strait (upper left) and Bosphorus Strait (lower right).

Model experiments presented focus on the period of the TSS08 sea trial (a joint project between the NATO Undersea Research Center and the U.S. Naval Research Laboratory), starting in late August and extending through November 2008. During one-way coupling as shown schematically in Figure

2, ADCIRC is initialized by temperature, salinity, velocity and water surface elevation fields from HYCOM-AMB solutions. At the open ocean boundaries, HYCOM-AMB values for elevation, temperature and salinity are updated daily throughout the ADCIRC model simulation. Surface forcing for both models is derived from the Navy's Coupled Ocean-Atmospheric Mesoscale Prediction System (COAMPS). The capability of ADCIRC to represent two-layer stratified flow dynamics both in the straits and in the Marmara Sea is examined along with the response of the currents and density structure over the water column to wind forcing. Observations include measured currents from ADCP moorings located at the ends of each strait in the TSS, CTD casts along the Dardanelles Strait and drifter deployments in the Marmara Sea. Observed features of interest include flow reversals in the straits during strong storm events, high-frequency current variability, and the persistence of circulation gyres in the Marmara Sea.




Fig. 2. The ADCIRC-HYCOM coupled model system.

## Acknowledgements

This work is supported by the Office of Naval Research under the 6.2 NRL program "Development of a Multi-Scale Coupled Ocean Model System: Application to the Turkish Straits". Computer time has been granted by the DoD High Performance Computing Modernization Program at NAVOCEANO. The authors thank Dr. Jeff Book for processing ADCPs collected during the TSS08 field campaign and Drs. A Birol Kara and A. J. Wallcraft for development of the HYCOM-AMB model. This paper is contribution number NRL/PP/7320-09-0094.

## References

- 1 - Bleck, R., 2002. An oceanic general circulation model framed in hybrid isopycnal-cartesian coordinates. *Ocean Model.*, 4: 55-88.
- 2 - Dresback K. M. and Kolar, R. L., 2009. Formulation and numerical implementation of the 3D Transport within the ADCIRC finite element code version 48.xx. *ADCIRC Theory Rep. (DRAFT)*.
- 3 - Kanarska, Y. and Maderich, V., 2008. Modelling of seasonal exchange flows through the Dardanelles Strait. *Estuar. Coast. Shelf S.*, 79: 449-458.
- 4 - Kourafalou, V. and Tsirlas, K., 2007. A nested circulation model for the North Aegean Sea. *Ocean Sci.*, 3: 1-16.
- 5 - Luettich Jr., R. A. and Westerink, J. J., 2004. Formulation and numerical implementation of the 2D/3D ADCIRC finite element model version 44.XX. *ADCIRC Theory Rep.*. Found at [www.adcirc.org](http://www.adcirc.org)
- 6 - Oguz, T., 2005. Hydraulic adjustments of the Bosphorus exchange flow. *Geophys. Res. Lett.*, 32: L06604, doi: 10.1029/2005GL022353.
- 7 - Staneva, J. V., Dietrich, D. E., Stanev, E. V., and Bowman, M. J., 2001. Rim current and coastal eddy mechanisms in an eddy-resolving Black Sea general circulation model. *J. Mar. Sys.*, 31: 137-157.